


## Burgess reagent

### Methoxycarbonylsulfamoyl triethylammonium hydroxide, inner salt

<b>CAS</b>	<b>29684-56-8</b>
<b>FW</b>	<b>238,30</b>
<b>Melting point</b>	<b>72-74 °C</b>



Synthetic Applications of *N*-Carboalkoxysulfamate Esters

Burgess, E.M.; Penton Jr., H.R.; Taylor, E.A.; *J. Am. Chem. Soc.*, 92 (1970) 5224;

Thermal reactions of alkyl *N*-carbomethoxysulfamate esters

Burgess, E.M.; Penton Jr., H.R.; Taylor, E.A.; *J. Org. Chem.*, 38 (1973) 26;

New synthetic reactions. Synthesis of cyclobutanes, cyclobutenes and cyclobutanones.

Applications in geminal alkylation

Trost, B.M.; Keeley, D.E.; Arndt, H.C.; Bogdanowicz, M.J.; *J. Am. Chem. Soc.*, 99 (1977) 3088;

Stereoselective thermal rearrangement of *syn*-7-(1,2-butadienyl)-1-methylbicyclo[2.2.1]hept-2-ene [*syn*-7-(3-methylallenyl)-1-methylnorbornene]

Duncan, J.A.; Hendricks, R.T.; Kwong, K.S.; *J. Am. Chem. Soc.*, 112 (1990) 8433;

Synthesis of a 1,9-dideoxy-forskolin derivative

Zimmermann, S.; Bick, S.; Welzel, P.; Meuer, H.; Sheldrick, W.S.; *Tetrahedron*, 51 (1995) 2947;

Enantioselective conjugate radical addition reactions mediated by Chiral Lewis acid complexes of (*R,R*)-4,6-dibenzofurandiyl-2,2'-bis(4-phenyloxazoline) (DBFOX/Ph)

Iserloh, U.; Curran, D.P.; Kanemasa, S.; *Tetrahedron:Asymmetry*, 10 (1999) 2417;

(Alkenyl)(amino)carbene complexes: potential starting materials for the synthesis of cyclopropylacetic acid derivatives

Papagni, A.; Maiorana, S.; Licandro, E.; Manzotti, R.; Baldoli, C.; *Eur. J. Org. Chem.*, (2001) 1149;

A novel regio- and stereoselective synthesis of sulfamidates from 1,2-diols using Burgess and related reagents: a facile entry into  $\beta$ -amino alcohols  
Nicolaou, K.C.; Huang X.; Snyder, S.A.; Rao, P.B.; Bella, M.; Reddy, M.V.; *Angew. Chem. Int. Ed.*, 41 (2002) 834;

A New method for the synthesis of nonsymmetrical sulfamides using Burgess-type Reagents  
Nicolaou, K.C.; Longbottom D.A.; Snyder, S.A.; Nalbandian A.Z.; Huang X.; *Angew. Chem. Int. Ed.*, 41 (2002) 3866;

Synthetic studies on thioStrepton family of peptide antibiotics: synthesis of the tetrasubstituted dihydroquinoline portion of siomycin D<sub>1</sub>  
Higashibayashi, S.; Mori, T.; Shinko, K.; Hashimoto, K.; Nakata, M.; *Heterocycles*, 57 (2002) 111;

Oxidation of benzoin to benzil using Burgess reagent  
Jose, B.; Unni, M.V.V.; Prathapan, S.; Vadakkan, J.J.; *Synt. Comm.*, 32 (2002) 2495;

New application of Burgess reagent in its reaction with epoxides  
Rinner, U.; Adams, D.R.; dos Santos, M.L.; Abboud, K.A.; Hudlicky, T.; *Synlett*, (2003) 1247;

Design, synthesis and evaluation of aza inhibitor of chorismate mutase  
Hediger, M.E.; *Bioorg. Med. Chem.*, 12 (2004) 4995;

New uses for the Burgess reagent in chemical synthesis: methods for the facile and stereoselective formation of sulfamidates, glycosylamines, and sulfamides.  
Nicolaou, K.C.; Snyder, S.A.; Longbottom D.A.; Nalbandian A.Z.; Huang X.; *Chem. Eur. J.*, 10 (2004) 5581;

A New method for the stereoselective synthesis of  $\alpha$ - and  $\beta$ -glycosylamines using the Burgess reagent  
Nicolaou, K.C.; Snyder, S.A.; Nalbandian A.Z.; Longbottom D.A.; *J. Am. Chem. Soc.*, 126 (2004) 6235;

One-pot easy conversion of Baylis-Hillman adducts into carbamates of unsaturated  $\beta$ -amino acids  
Mamaghani, M.; Badrian, A.; *Tetrahedron Letters*, 45 (2004) 1547;

A convenient enantioselective synthesis of (*S*)- $\alpha$ -trifluoromethylisoserine  
Avenoza, A.; Busto, J.H.; Jimenez-Oses, G.; Peregrina, J.M.; *J. Org. Chem.*, 70 (2005) 5721;

*De novo* enantioselective syntheses of galacto-sugars and deoxy sugars via the iterative dihydroxylation of dienoate  
Ahmed, Md.M.; Berry, B.P.; Hunter, T.J.; Tomcik, D.J.; O'Doherty, G.A.; *Org. Lett.*, 7 (2005) 745;

Activation of carboxylic acid by Burgess reagent: an efficient route to acyl ureas and amides  
Wodka, D.; Robbins, M.; Lan, P.; Martin, R.L.; Athanasopoulos, J.; Makara, G.M.; *Tetrahedron Letters*, 47 (2006) 1547;